

Pharmacologic and Non-Pharmacologic Methods in the Prevention of HIV: 2023

Christopher W. Blackwell, Ph.D., APRN,
ANP-BC, AGACNP-BC, CNE, FAANP,
FAAN

*Associate Professor & Program Director
AGACNP Programs, College of Nursing
Academic Health Sciences Center
University of Central Florida
Orlando, Florida*



DISCLOSURES

Dr. Blackwell has no conflicts of interest or other disclosures for this presentation.



Images and Figures: All images and figures obtained through Creative Commons license or through the CDC. Note, the CDC openly encourages use of its data and free and open use of its data tables and figures (see <https://www.cdc.gov/other/agencymaterials.html>).

Image licensed through Creative Commons: <https://www.picpedia.org/real-estate/d/disclosures.html>



NEW ORLEANS ERNEST N. MORIAL
CONVENTION CENTER



OBJECTIVES

1. At the end of this presentation, participants will outline most recent epidemiology of HIV infection among adults and adolescents in the United States.
2. At the end of this presentation, participants will identify two FDA-approved oral pharmacologic agents and one injectable pharmacologic agent used to prevent HIV infection using pre-exposure prophylaxis (PrEP).
3. At the end of this presentation, participants will describe the pharmacologic prevention of HIV through post-exposure prophylaxis (PEP).
4. At the end of this presentation, participants will acknowledge the protective effect of undetectable HIV RNA levels (viral load) in eliminating sexual transmission of HIV among high-risk groups.
5. At the end of this presentation, participants will identify the responsibilities of nurse practitioners in leading future scholarship and clinical practice in preventing HIV.




Incidence of HIV Infections & AIDS

- Review of Centers for Disease Control and Prevention (CDC) Data: Updated through 2020 (2016-2020)
- These can all be obtained from the *2020 CDC HIV Surveillance Report, 33*:
- <https://www.cdc.gov/hiv/pdf/library/reports/surveillance/cdc-hiv-surveillance-report-2020-updated-vol-33.pdf>
- The figures on slides 4-11 all come from these CDC sources



Incidence of HIV Infection & AIDS 2016-2020: Take Away Points (2020)

Figure 1. Rates of diagnoses of HIV infection among persons aged ≥13 years, 2020 (COVID-19 pandemic)—United States and 6 dependent areas




Note. Data for 2020 should be interpreted with caution due to the impact of the COVID-19 pandemic on access to HIV testing, care-related services, and case surveillance activities in state/local jurisdictions. Asterisk (*) indicates incomplete reporting.



Incidence of HIV Infection & AIDS 2016-2020: Take Away Points (2020)

Ir

Figure 2. Percentages of diagnoses of HIV infection among persons aged ≥ 13 years, by gender, 2020 (COVID-19 pandemic)—United States and 6 dependent areas




Note. Data for 2020 should be interpreted with caution due to the impact of the COVID-19 pandemic on access to HIV testing, care-related services, and case surveillance activities in state/local jurisdictions.

- **Age group** (Figure 3):
 - 13–24 years: 20%
 - 25–34 years: 37%
 - 35–44 years: 20%
 - 45–54 years: 13%
 - ≥ 55 years: 10%




Incidence of HIV Infection & AIDS 2016-2020: Take Away Points (2020)

Figure 3. Percentages of diagnoses of HIV infection among persons aged ≥13 years, by age at diagnosis, 2020 (COVID-19 pandemic)—United States and 6 dependent areas




Note. Data for 2020 should be interpreted with caution due to the impact of the COVID-19 pandemic on access to HIV testing, care-related services, and case surveillance activities in state/local jurisdictions.



Incidence of HIV Infection & AIDS 2016-2020: Take Away Points (2020)

Figure 5. Percentages of diagnoses of HIV infection among persons aged ≥ 13 years, by race/ethnicity, 2020 (COVID-19 pandemic)—United States and 6 dependent areas




Note. Data for 2020 should be interpreted with caution due to the impact of the COVID-19 pandemic on access to HIV testing, care-related services, and case surveillance activities in state/local jurisdictions.

^a Hispanic/Latino persons can be of any race.



Incidence of HIV Infection & AIDS 2016-2020: Take Away Points (2020)

Figure 6. Percentages of diagnoses of HIV infection among persons aged ≥13 years, by sex assigned at birth and transmission category, 2020 (COVID-19 pandemic)—United States and 6 dependent areas




Note. Data have been statistically adjusted to account for missing transmission category. See section D4 in Technical Notes for more information on transmission categories.



Incidence of HIV Infection & AIDS 2016-2020: Take Away Points (2020)

Figure 13. Percentages of diagnoses of HIV infection among men who have sex with men, by age group and race/ethnicity, 2020 (COVID-19 pandemic)—United States and 6 dependent areas



Note. Data have been statistically adjusted to account for missing transmission category. Data for 2020 should be interpreted with caution due to the impact of the COVID-19 pandemic on access to HIV testing, care-related services, and case surveillance activities in state/local jurisdictions.


^a Hispanic/Latino persons can be of any race.



Incidence of HIV Infection & AIDS: 2015-2019 Take Away Points

Ir

Figure 22. Percentages of diagnoses of HIV infection and population among females aged ≥13 years, by race/ethnicity, 2020 (COVID-19 pandemic)—United States



Note. Data for 2020 should be interpreted with caution due to the impact of the COVID-19 pandemic on access to HIV testing, care-related services, and case surveillance activities in state/local jurisdictions.

^a Hispanic/Latino persons can be of any race.



HIV REPLICATION CYCLE




Image licensed through Creative Commons by NIH NIAID:<https://www.niaid.nih.gov/diseases-conditions/hiv-replication-cycle>



Pathophysiology of HIV Infection

- HIV is a retrovirus, transcribing RNA-containing genetic material into DNA of the host cell nucleus by using an enzyme called reverse transcriptase
- Glycoproteins allow HIV to attach to CD4 Cell and incorporate its RNA into the cell membrane, which then transcribes the RNA to DNA using reverse transcriptase
- This is then integrated into the CD4 nucleus using integrase. Integrated viral genes then transcribe back into genomic RNA and messenger RNA, which are translated to viral proteins
- These proteins then are cleaved with protease into new HIV particles, which release to infect other cells
- HIV progresses to AIDS
- Seroconversion (HIV- → HIV+) typically occurs in 2-12 weeks post-exposure. 95% (1 month; 99.9% by week 12)



Pathophysiology of HIV Infection

- After seroconversion, HIV antibody titers decrease as infected cells are sequestered in the lymph nodes
- This is the latent period, lasting up to 10 years
- During this period, CD4 cell lines drop as a result of infection and lysis of healthy T-Helper cells



Pathophysiology of HIV Infection

- As CD4 cells continue to decline, the patient becomes susceptible to opportunistic infections, malignancies, and neurological diseases
 - AIDS develops
- A very few HIV+ individuals are termed “Non-Progressors”



Pathogenic Process of HIV/AIDS

- Exposure to HIV
- HIV Infection
- Seroconversion
- Latency Period
- Initial Symptoms of Immunodeficiency and Declining Immune Function
- Immune System Failure and AIDS
- Severe Immune Deficiency



Pathogenic Process of HIV

- Important Points:
- Transmission of HIV is possible at any stage of the disease process
- Risk to health workers is overall small
- With blood product screening emerging in 1985, transfusion-related HIV transmission decreased dramatically
- Since the introduction of maternal antiretroviral therapy, HIV transmission from mom to child has decreased
- *Practically Preventable*



Prevention of HIV Transmission

- Sexual Transmission:
 - Alteration in Sexual Behaviors
 - Women more susceptible via vaginal mucosa compared to male penis
 - Anal intercourse (regardless of orientation) also risky secondary to rectal trauma, tearing, and fistula formation
 - Oral sex is actually very low risk
 - Viral Load is NOT a determinant of degree of safeness (theoretically)—CDC (2017) issued newer statement about this



Prevention of HIV Transmission

- Pharmacologic: PrEP and PEP
- Parenteral Transmission:
 - Proper cleaning of drug paraphernalia:
 - Fill with water (tap to loosen blood debris) and flush →
Fill with bleach and then shake for 30 seconds, flush →
Repeat x 3 →
Fill with water, shake and tap x 30 seconds, flush →
Repeat x 3
 - Participation in needle exchange programs

Image licensed through Creative Commons by PXhere: <https://pxhere.com/en/photo/774066>



Prevention of HIV Transmission

- Perinatal Transmission:
 - HIV transmission thought to occur transplacentally in utero, intrapartally during exposure to blood and vaginal secretions during childbirth, or postpartally through breast milk



Prevention of HIV Transmission

- Perinatal Transmission (Ctd):
- Review of prior HIV-related illnesses and past CD4 T lymphocyte (CD4) cell counts and plasma HIV RNA levels;
- Current CD4 cell count;
- Current plasma HIV RNA copy number;
- Assessment of the need for prophylaxis against opportunistic infections such as *Pneumocystis jirovecii* pneumonia and *Mycobacterium avium* complex (see Adult and Adolescent Opportunistic Infections Guidelines)



Prevention of HIV Transmission

- Perinatal Transmission (Ctd):
- Screening for hepatitis C virus and tuberculosis in addition to standard screening for hepatitis B virus (HBV) infection;
- Assessment of the need for immunizations per guidelines from the American College of Obstetricians and Gynecologists, with particular attention to hepatitis A, HBV, influenza, pneumococcus, and Tdap immunizations;
- Complete blood cell count and renal and liver function testing;
- HLA-B*5701 testing if abacavir (Ziagen®) use is anticipated;
- History of prior and current antiretroviral (ARV) drug use, including prior ARV use for prevention of perinatal transmission or treatment of HIV and history of adherence problems



Prevention of HIV Transmission

- Perinatal Transmission (Ctd):
 - Infected with HIV and on ART?:
 - Keep taking ART!
 - Infected with HIV and not on ART or with unknown or high HIV RNA load?:
 - Begin zidovudine (Retrovir®) IV near time of delivery
 - C-section in @ 38 weeks gestation
 - Neonate will also be treated with ART
 - Most recent guidelines (updated 2023):
<https://clinicalinfo.hiv.gov/en/guidelines/perinatal/antiretroviral-management-newborns-perinatal-hiv-exposure-or-hiv-infection>



Screening for HIV

- ELISA → Western Blot (99.5% accurate)
 - Newer guidelines (CDC, 2014) are calling for substitution of Western Blot with antigen tests that differentiate HIV1 from HIV2.
- Confidentiality is **INCREDIBLY** important (i.e., no phone messages, personal names, etc.).
- Pre-Test and Post-Test Counseling can be valuable but is NOT CDC recommended as a requirement any longer.
 - Check your state regulations for guidance
- General consent for Tx implies consent for HIV



Image licensed through Creative Commons by Wallpaper Flare: <https://www.wallpaperflare.com/the-hand-child-cover-up-do-not-do-not-shoot-the-prohibition-of-wallpaper-atdyk>



Screening for HIV

- Antibody tests are specifically designed for the routine testing of HIV in adults, are inexpensive, and are very accurate
- Antibody tests give false negatives results during the *window period* of between three weeks and six months from the time of HIV infection until the immune system produces detectable amounts of antibodies
- Much screening done as POS
 - e.g., OraSure® OraQuick® testing methods




Image licensed through Creative Commons by Wikimedia Commons: <https://commons.wikimedia.org/wiki/File:Oraquick.jpg>



Screening for HIV

- Most people have detectable antibodies after three months
- A six-month window is extremely rare with modern antibody testing
- During this window period an infected person can transmit HIV to others, without their HIV infection being detectable using an antibody test
- ART during the window period can delay the formation of antibodies and extend the window period beyond 12 months



Screening for HIV

- The specificity of rapid antibody tests in low-risk populations has not been evaluated
- Designed for high-risk individuals
 - OraSure® saliva—collected on oral wand device placed between gum and cheek for 2-5 min--mixed in a vial with solution, wand snapped off, vial closed and sent to lab
 - It is an antibody test which first employs ELISA, then Western Blot
 - OraQuick® Advance is an HIV test which uses saliva, plasma, fingerstick, or whole blood specimen
 - Sample is obtained and mixed in a buffer → Device inserted into buffer → Results in 20-40 min
 - CLIA-waived for saliva, fingerstick, and venipuncture whole blood
 - There is also a urine test; it employs both the ELISA and the Western Blot method
 - Home Access Express HIV-1 Test is a FDA-approved home test: the patient collects a drop of blood and mails the sample to a laboratory; the results are obtained over the phone



Screening for HIV

- **Antigen Tests:**

- The **p24 antigen test** detects the presence of the p24 protein of HIV (also known as CA), a major core protein of the virus
- This test is now used routinely to screen blood donations, thus reducing the window to about 16 days

- **Nucleic Acid-Based Tests:**

- Nucleic acid-based tests amplify and detect a 142 base target sequence located in a highly conserved region of the HIV *gag* gene
- Since 2001, donated blood in the US has been screened with nucleic acid-based tests, shortening the window to about 12 days
- Since these tests are relatively expensive, the blood is screened by first pooling some 10-20 samples, testing these together, and if the pool tests positive, each sample is retested individually



HIV/ AIDS Surveillance and Dx

- CD4 Testing:
 - Declining CD4 T-cell counts are a marker of the progression of HIV infection.
 - In HIV+ people, AIDS is officially diagnosed when the count drops below 200 cells or when certain opportunistic infections occur; CDC guidelines recommend beginning ART AT TIME OF Dx (2015)
 - Low CD4 T-cell counts are associated with a variety of conditions, including many viral infections, bacterial infections, parasitic infections, sepsis, tuberculosis, coccidioidomycosis, burns, trauma, intravenous injections of foreign proteins, malnutrition, over-exercising, pregnancy, normal daily variation, psychological stress, and social isolation



HIV/ AIDS Surveillance and Dx

- CD4 Testing:
 - The lower the number of T cells, the lower the immune system's function will be
 - Normal T4 counts are between 500 and 1500 CD4+ T cells per microliter and the counts may fluctuate in healthy people, depending on recent infection status, nutrition, exercise and other factors -- even the time of day
 - Women tend to have somewhat lower counts than men



HIV/ AIDS Surveillance and Dx

- Viral Load Testing:
 - Evidence shows that keeping the viral load levels as low as possible for as long as possible decreases the complications of HIV disease and prolongs life
 - Most recent public health guidelines state that treatment should be considered for asymptomatic HIV-infected people AT TIME OF Dx
 - There are several methods for testing viral load; results are not interchangeable, so it is important that the same method be used each time
 - Keep viral loads undetectable = decrease/ eliminate transmission



Prophylactic Prevention of HIV Infection: Post-Exposure

- Although large-scale studies about PEP are lacking, PEP is clinically effective (80%) and recommended (Landovitz & Currier, 2009) when:
 - The source is known to be HIV+
 - The source is of unknown serostatus (test source in occupational exposure)
 - The source has an increased likelihood of being HIV+:
 - MSM, MSM/W, commercial sex workers, history of incarceration, residence in a county with a seroprevalence rate \geq 1%
 - The behavior has an increased (\geq 1%) likelihood of transmitting HIV:
 - Receptive Anal Intercourse = 1%-30% chance of infection
 - Insertive Anal Intercourse = .1-10% chance of infection
 - Receptive Vaginal Intercourse = .1-10% chance of infection
 - Insertive Vaginal Intercourse = .1-1% chance of infection
 - Oral Intercourse: Few documented cases
 - Needle Sharing: .67% per needle-sharing event
- Ideally, begin PEP within 36 hours but no more than 72 hours after exposure



Prophylactic Prevention of HIV Infection: Post-Exposure

Table 1. Estimated per-act risk for acquiring human immunodeficiency virus (HIV) from an infected source, by exposure act^a

Exposure type	Rate for HIV acquisition per 10,000 exposures
Parenteral	
Blood transfusion	9,250
Needle sharing during injection drug use	63
Percutaneous (needlestick)	23
Sexual	
Receptive anal intercourse	138
Receptive penile-vaginal intercourse	8
Insertive anal intercourse	11
Insertive penile-vaginal intercourse	4
Receptive oral intercourse	Low
Insertive oral intercourse	Low
Other^b	
Biting	Negligible
Spitting	Negligible
Throwing body fluids (including semen or saliva)	Negligible
Sharing sex toys	Negligible

Source: <http://www.cdc.gov/hiv/policies/law/risk.html>


^a Factors that may increase the risk of HIV transmission include sexually transmitted diseases, acute and late-stage HIV infection, and high viral load. Factors that may decrease the risk include condom use, male circumcision, antiretroviral treatment, and preexposure prophylaxis. None of these factors are accounted for in the estimates presented in the table.

^b HIV transmission through these exposure routes is technically possible but unlikely and not well documented.

Source: <https://stacks.cdc.gov/view/cdc/38856>



Prophylactic Prevention of HIV Infection: Post-Exposure



Source: <https://stacks.cdc.gov/view/cdc/38856>



Prophylactic Prevention of HIV Infection: Post-Exposure

Table 5. Preferred and alternative antiretroviral medication 28-day regimens for nPEP^{a,b}

Age group	Preferred/ alternative	Medication
Adults and adolescents aged ≥ 13 years, including pregnant women, with normal renal function (creatinine clearance ≥ 60 mL/min)	Preferred	A 3-drug regimen consisting of tenofovir DF 300 mg and fixed dose combination emtricitabine 200 mg (Truvada ^c) once daily with raltegravir 400 mg twice daily or dolutegravir 50 mg once daily
	Alternative	A 3-drug regimen consisting of tenofovir DF 300 mg and fixed dose combination emtricitabine 200 mg (Truvada) once daily with darunavir 800 mg (as 2, 400-mg tablets) once daily and ritonavir ^b 100 mg once daily
Adults and adolescents aged ≥ 13 years with renal dysfunction (creatinine clearance ≤ 59 mL/min)	Preferred	A 3-drug regimen consisting of zidovudine and lamivudine, with both doses adjusted to degree of renal function with raltegravir 400 mg twice daily or dolutegravir 50 mg once daily
	Alternative	A 3-drug regimen consisting of zidovudine and lamivudine, with both doses adjusted to degree of renal function with darunavir 800 mg (as 2, 400-mg tablets) once daily and ritonavir ^b 100 mg once daily

Source: <https://stacks.cdc.gov/view/cdc/38856>



Prophylactic Prevention of HIV Infection: Post-Exposure

Table 2. Recommended schedule of laboratory evaluations of source and exposed persons for providing nPEP with preferred regimens

Test	Source	Exposed persons			
	Baseline	Baseline	4–6 weeks after exposure	3 months after exposure	6 months after exposure
	For all persons considered for or prescribed nPEP for any exposure				
HIV Ag/Ab testing ^a (or antibody testing if Ag/Ab test unavailable)	✓	✓	✓	✓	✓ ^b
Hepatitis B serology, including: hepatitis B surface antigen hepatitis B surface antibody hepatitis B core antibody	✓	✓	—	—	✓ ^c
Hepatitis C antibody test	✓	✓	—	—	✓ ^d
	For all persons considered for or prescribed nPEP for sexual exposure				
Syphilis serology ^e	✓	✓	✓	—	✓
Gonorrhea ^f	✓	✓	✓ ^g	—	—
Chlamydia ^f	✓	✓	✓ ^g	—	—
Pregnancy ^h	—	✓	✓	—	—
	For persons prescribed tenofovir DF+ emtricitabine + raltegravir or tenofovir DF+ emtricitabine + dolutegravir				
Serum creatinine (for calculating estimated creatinine clearance ⁱ)	✓	✓	✓	—	—
Alanine transaminase, aspartate aminotransferase	✓	✓	✓	—	—
	For all persons with HIV infection confirmed at any visit				
HIV viral load	✓	✓	✓	✓ ^j	✓
HIV genotypic resistance	✓	✓	✓	✓ ^j	✓

Abbreviations: Ag/Ab, antigen/antibody combination test; HIV, human immunodeficiency virus; nPEP, nonoccupational postexposure prophylaxis; tenofovir DF, tenofovir disoproxil fumarate.

^a Any positive or indeterminate HIV antibody test should undergo confirmatory testing of HIV infection status.

^b Only if hepatitis B infection was acquired during the original exposure; delayed HIV seroconversion has been seen in persons who simultaneously acquire HIV and hepatitis B infection.

^c If exposed person susceptible to hepatitis B at baseline.

^d If exposed person susceptible to hepatitis C at baseline.

^e If determined to be infected with syphilis and treated, should undergo serologic syphilis testing 6 months after treatment.

^f Testing for chlamydia and gonorrhea should be performed using nucleic acid amplification tests. For patients diagnosed with a chlamydia or gonorrhea infection, retesting 3 months after treatment is recommended.

- For men reporting insertive vaginal, anal, or oral sex, a urine specimen should be tested for chlamydia and gonorrhea.
- For women reporting receptive vaginal sex, a vaginal (preferred) or endocervical swab or urine specimen should be tested for chlamydia and gonorrhea.
- For men and women reporting receptive anal sex, a rectal swab specimen should be tested for chlamydia and gonorrhea.
- For men and women reporting receptive oral sex, an oropharyngeal swab should be tested for gonorrhea. (<http://www.cdc.gov/std/tg2015/tg-2015-print.pdf>)

^g If not provided presumptive treatment at baseline, or if symptomatic at follow-up visit.

^h If woman of reproductive age, not using effective contraception, and with vaginal exposure to semen.

ⁱ eCrCl = estimated creatinine clearance calculated by the Cockcroft-Gault formula; eCrClCG = [(140 – age) x ideal body weight] + (serum creatinine x 72) (x 0.85 for females).

^j At first visit where determined to have HIV infection.

Source: <https://stacks.cdc.gov/view/cdc/38856>



Pre-Exposure Prophylaxis


- PrEP therapy is indicated for patients considered high risk for sexually acquired HIV
- Examples of such individuals might include a non-HIV-infected partner of an HIV infected individual



Image licensed through Creative Commons by PXhere: <https://pxhere.com/en/photo/1444649>



Pre-Exposure Prophylaxis



Preexposure Prophylaxis for the Prevention of HIV Infection in the United States – 2021 Update Clinical Practice Guideline

Source: <https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2021.pdf>



Oral Pre-Exposure Prophylaxis

- The PrEP dosage is one tablet (emtricitabine 200 mg and tenofovir disoproxil fumarate 300 mg [FTC/TDF {Truvada®}] *or* emtricitabine 200 mg and tenofovir alafenamide 25 mg [FTC/TAF {Descovy®}])
- Emtricitabine 200 mg and tenofovir disoproxil fumarate 300 mg
 - *Approved for adult and adolescent cisgender/ transgender males and females*
- Emtricitabine 200 mg and tenofovir alafenamide 25 mg
 - *Approved for adult and adolescent cisgender males and transgender females only*
- Taken PO with or without food and should be prescribed with a frequency of once daily
- In addition to the medication, which should be prescribed in no more than a 90-day supply, the patient should be educated about risk reduction strategies, particularly consistent use of condoms during every sexual encounter



Oral Pre-Exposure Prophylaxis

Treatment Monitoring Recommendations

Test	Screening/Baseline Visit	Q 3 months	Q 6 months	Q 12 months	When stopping PrEP
HIV Test	X*	X			X*
eCrCl	X		If age ≥ 50 or eCrCL < 90 ml/min at PrEP initiation	If age < 50 and eCrCl ≥ 90 ml/min at PrEP initiation	X
Syphilis	X	MSM /TGW	X		MSM/TGW
Gonorrhea	X	MSM /TGW	X		MSM /TGW
Chlamydia	X	MSM /TGW	X		MSM /TGW
Lipid panel (F/TAF)	X			X	
Hep B serology	X				
Hep C serology	MSM, TGW, and PWID only			MSM, TGW, and PWID only	

* Assess for acute HIV infection (see Figure 4)

- In females, document a negative urine pregnancy test


Source: <https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2021.pdf>



Oral Pre-Exposure Prophylaxis

- “2-1-1” Dosing

Figure 8 Schedule for “2-1-1” Dosing



Based on the timing of subsequent sexual events, MSM should be instructed to take additional doses as follows:

- If sex occurs on the consecutive day after completing the 2-1-1 doses, take 1 pill per day until 48 hours after the last sexual event.
- If a gap of <7 days occurs between the last pill and the next sexual event, resume 1 pill daily.
- If a gap of ≥ 7 days occurs between the last pill and next sexual event, start again with 2 pills.



Oral Pre-Exposure Prophylaxis



- Prevention effectiveness occurs after 7-20 days
- The financing of antiretrovirals for PrEP is emerging as an important healthcare policy issue
- Daily cost of Oral PrEP up to \$1750/month, which equals \$21,000 per year
- Additional monitoring and screening costs per person have been estimated to be \$1,300 per year.
- USPTF A Recommendation (6/19), thus private insurance companies must cover PrEP

Image licensed through Creative Commons by flickr: https://www.flickr.com/photos/tracy_olson/61056391



NEW ORLEANS ERNEST N. MORIAL
CONVENTION CENTER

Injectable Pre-Exposure Prophylaxis

- Data strongly suggest use of every 2-month injectable cabotegravir (Apretude®) (600 mg/ 3mL), in concert with safer sex practices, reduces risk of HIV-1 acquisition by approximately 99%, with only a very small number of infections (related to resistant HIV strains) occurring in those who are adherent
- This administration approach might be beneficial to those at risk for HIV infection for which adherence may be challenging
 - Young men who have sex with men (MSM), people with substance abuse disorders, those living in poverty or who have depression, conceal their use of PrEP, or are otherwise challenged with adherence
- Before starting injectable PrEP, patients should be screened for HIV, bacterial STIs, and hepatitis B; baseline renal and hepatic function should also be assessed
 - MSM and people who injecting drugs (PWIDs) should also be screened for hepatitis C
 - A negative HIV screening should be obtained within 1 week of starting injectable cabotegravir



Injectable Pre-Exposure Prophylaxis

- Using an injectable lead-in strategy, an initial dose of cabotegravir 600 mg (in 3 mL) is administered IM in the dorsal gluteal muscle; a second dose is given 4-weeks after this first dose; then every 8 weeks thereafter.
 - An oral lead-in approach uses this same schedule but is preceded by at least 28 days of daily oral cabotegravir 30 mg, with the first injectable dose given on the final day of the oral lead-in.
- Injectable PrEP patients should be screened for HIV 1-month after the first injection and at follow-up visits every 2 months (both HIV Ag/Ab test AND HIV-1 RNA assay); MSM and transgender women who have sex with men should additionally be screened for bacterial STIs (oral, rectal, urine, blood) every 4 months while sexually active heterosexual men and women and men (vaginal, rectal, urine—as indicated) every 6 months
- Ongoing lipid and renal evaluations are unnecessary.
- Common adverse events associated with injectable cabotegravir include injection site reactions, diarrhea, headache, pyrexia, and fatigue.
- Clinicians should provide continuing guidance on safer sexual decision making and answer any questions that arise while using PrEP throughout the ongoing regimen.



Pre-Exposure Prophylaxis

- Evaluating patient appropriateness for PrEP, performing pretreatment evaluations prior to initiation of treatment, and close monitoring of therapy are all responsibilities NPs will assume as this treatment becomes more widespread in the U.S. healthcare system.
- Cost of the therapy is also a major blockade to its implementation, and this will continue to be a prevalent issue in the foreseeable future



Pre-Exposure Prophylaxis AANP Resources

- Two Practice Briefs are available:
 - Log into AANP Site
 - Click on Practice
 - Click on Clinical Resources
 - Scroll down to Point of Care Tools and Clinical Practice Briefs
 - Click Access Point of Care Tools
 - Click on Clinical Practice Briefs
 - Scroll down to Infectious Disease
 - Click on “View the Brief” underL
 - Pre-Exposure Prophylaxis for HIV Prevention– Injectable
 - Pre-Exposure Prophylaxis for HIV Prevention– Oral



Image licensed through Creative Commons by Wikimedia Commons: https://commons.wikimedia.org/wiki/File:American_Association_of_Nurse_Practitioners_logo_031918.png



Moving Forward

- Community and Public Health Outreach
- Prevention Education in the Clinical Setting
- Future Research and the Responsibility of the Nurse Practitioner




Image licensed through Creative Commons by Pixy: <https://pixy.org/1601486/>



References

Please see the supplemental handout, which includes a bibliography and additional resources for more information.

Scan the QR Code to access the online bibliography!



Bibliography Compiled by:

Andrew Todd, MLIS, BSN, RN

Reference Librarian, College of Nursing

University of Central Florida Libraries



References

- Althoff, K. N., Stewart, C. N., Humes, E., Zhang, J., Gerace, L., Boyd, C. M., Wong, C., Justice, A. C., Gebo, K. A., Thorne, J. E., Rubtsova, A. A., Horberg, M. A., Silverberg, M. J., Leng, S. X., Rebeiro, P. F., Moore, R. D., Buchacz, K., & Kasaie, P. (2022). The shifting age distribution of people with HIV using antiretroviral therapy in the United States. *AIDS (02699370)*, *36*(3), 459–471. <https://doi.org/10.1097/QAD.0000000000003128>
- Beer, L., Tie, Y., McCree, D. H., Demeke, H. B., Marcus, R., Padilla, M., Khalil, G., & Shouse, R. L. (2022). HIV Stigma among a national probability sample of adults with diagnosed HIV-United States, 2018-2019. *AIDS and Behavior*, *26*(Suppl 1), 39–50. <https://doi.org/10.1007/s10461-021-03414-6>
- Begley, R., Das, M., Zhong, L., Ling, J., Kearney, B. P., & Custodio, J. M. (2018). Pharmacokinetics of tenofovir alafenamide when coadministered with other hiv antiretrovirals. *Journal of Acquired Immune Deficiency Syndromes (1999)*, *78*(4), 465–472. <https://doi.org/10.1097/QAI.0000000000001699>
- Blackwell, C. W. (2008). Men who have sex with men and recruit bareback sex partners on the Internet: Implications for STI and HIV prevention and client education. *American Journal of Men's Health*, *2*(4), 306–313. <https://doi.org/10.1177/1557988307306045>
- Blackwell, C. W. (2014). Preexposure prophylaxis: An emerging clinical approach to preventing HIV in high-risk adults. *Nurse Practitioner*, *39*(9), 50–53. <https://doi.org/10.1097/01.NPR.0000452976.92052.fa>
- Blackwell, C. W. (2018). Preventing HIV Infection in high-risk adolescents using preexposure prophylaxis (PrEP). *The Journal of the Association of Nurses in AIDS Care : JANAC*, *29*(5), 770–774. <https://doi.org/10.1016/j.jana.2018.06.001>
- Blackwell, C. W., & Castillo, H. L. (2021). Human immunodeficiency virus pre-exposure prophylaxis: Use of emtricitabine/tenofovir alafenamide. *Journal for Nurse Practitioners*, *17*(6), 673–676. <https://doi.org/10.1016/j.nurpra.2021.01.002>
- Blackwell, C.W., & Lopez Castillo, H. (2022). Injectable cabotegravir: A new approach to HIV pre-exposure prophylaxis. *JNP-- The Journal for Nurse Practitioners*, *18*(9), 947-950.
- Caceres, B. A., Travers, J., Primiano, J. E., Luscombe, R. E., & Dorsen, C. (2020). Provider and LGBT Individuals' Perspectives on LGBT issues in long-term care: A Systematic Review. *The Gerontologist*, *60*(3), e169–e183. <https://doi.org/10.1093/geront/gnz012>



- Caughey, A. B., Krist, A. H., Wolff, T. A., Barry, M. J., Henderson, J. T., Owens, D. K., Davidson, K. W., Simon, M. A., & Mangione, C. M. (2021). USPSTF approach to addressing sex and gender when making recommendations for clinical preventive services. *JAMA: Journal of the American Medical Association*, 326(19), 1953–1961. <https://doi.org/10.1001/jama.2021.15731>
- Centers for Disease Control and Prevention (2023). *HIV Surveillance Report, 2020* (Volume 33). <https://www.cdc.gov/hiv/pdf/library/reports/surveillance/cdc-hiv-surveillance-report-2020-updated-vol-33.pdf>
- Crim, S. M., Yunfeng Tie, Beer, L., Weiser, J., Dasgupta, S., & Tie, Y. (2020). Barriers to antiretroviral therapy adherence among hiv-positive Hispanic and Latino men who have sex with men - United States, 2015-2019. *MMWR: Morbidity & Mortality Weekly Report*, 69(40), 1438–1442. <https://doi.org/10.15585/mmwr.mm6940a1>
- Crooks, E. T., Almanza, F., D’Addabbo, A., Duggan, E., Zhang, J., Wagh, K., Mou, H., Allen, J. D., Thomas, A., Osawa, K., Korber, B. T., Tsybovsky, Y., Cale, E., Nolan, J., Crispin, M., Verkoczy, L. K., & Binley, J. M. (2021). Engineering well-expressed, V2-immunofocusing HIV-1 envelope glycoprotein membrane trimers for use in heterologous prime-boost vaccine regimens. *PLoS Pathogens*, 17(10), 1–39. <https://doi.org/10.1371/journal.ppat.1009807>
- Crowe, S., Bennett, B., & Fordan, S. (2022). Impact of the 2014 CDC HIV testing guidelines on detection of acute HIV infections. *Journal of Clinical Virology : The Official Publication of the Pan American Society for Clinical Virology*, 146, 105058. <https://doi.org/10.1016/j.jcv.2021.105058>
- Dahlen, S., Connolly, D., Arif, I., Junejo, M. H., Bewley, S., & Meads, C. (2021). International clinical practice guidelines for gender minority/trans people: Systematic review and quality assessment. *BMJ Open*, 11(4), e048943. <https://doi.org/10.1136/bmjopen-2021-048943>
- Dunkle, L. M., Kotloff, K. L., Gay, C. L., Áñez, G., Adelglass, J. M., Barrat Hernández, A. Q., Harper, W. L., Duncanson, D. M., McArthur, M. A., Florescu, D. F., McClelland, R. S., Garcia-Fragoso, V., Riesenber, R. A., Musante, D. B., Fried, D. L., Safirstein, B. E., McKenzie, M., Jeanfreau, R. J., Kingsley, J. K., ... Dubovsky, F. (2022). Efficacy and safety of NVX-CoV2373 in Adults in the United States and Mexico. *The New England Journal of Medicine*, 386(6), 531–543. <https://doi.org/10.1056/NEJMoa2116185>
- Finlayson, T., Susan Cha, Ming Xia, Trujillo, L., Denson, D., Prejean, J., Kanny, D., Wejnert, C., Cha, S., & Xia, M. (2019). Changes in HIV preexposure prophylaxis awareness and use among men who have sex with men - 20 urban areas, 2014 and 2017. *MMWR: Morbidity & Mortality Weekly Report*, 68(27), 597–603. <https://doi.org/10.15585/mmwr.mm6827a1>
- Gilead 701*. (n.d.). Drugs.com. Retrieved March 11, 2022 from <https://www.drugs.com/imprints/gilead-701-9326.html>
- Gilead Sciences. (2022, January). *Highlights of prescribing information*. https://www.gilead.com/~media/Files/pdfs/medicines/hiv/descovy/descovy_pi.pdf.
- Gilead Sciences. (n.d.-a). *Simple dosing with Descovy for PrEP*. https://www.descovyhcp.com/dosing-and-product-offerings?utm_medium=cpc&utm_campaign=USA_GO_SEM_B_EX_Descovy-HCP-Stay+On+a+Gilead+Medication+Lead-Standard&utm_content=Descovy_Prescribing&utm_term=Descovy+prescribing+information&utm_source=google&gclid=EA1aIQobChMIu-j9zer47AIVkueGCh1fIA1pEAAAYASAAEglg3PD_BwE&gclid=aw.ds.



Gilead Sciences. (n.d.-b). *Co-pay support for the commercially insured*. <https://www.gileadadvancingaccess.com/hcp/financial-assistance/copay-support>.

Gilead Sciences. (n.d.-c). *Financial support for the uninsured—Multiple options available for enrollment*. <https://www.gileadadvancingaccess.com/hcp/financial-assistance/uninsured-support>.

Gilead Sciences. (n.d.-d). Understanding coverage options for patients. <https://www.gileadadvancingaccess.com/hcp/insurance>. Published 2020. Accessed on November 11, 2020.

Givens, M., Levison, J., & Rahangdale, L. (2021). Considerations and recommendations for pregnancy and postpartum care for people living with human immunodeficiency virus. *Obstetrics & Gynecology*, 138(1), 119–130. <https://doi.org/10.1097/AOG.0000000000004441>

Glidden, D. V., Das, M., Dunn, D. T., Ebrahimi, R., Zhao, Y., Stirrup, O. T., Baeten, J. M., & Anderson, P. L. (2021). Using the adherence-efficacy relationship of emtricitabine and tenofovir disoproxil fumarate to calculate background HIV incidence: A secondary analysis of a randomized, controlled trial. *Journal of the International AIDS Society*, 24(5), e25744. <https://doi.org/10.1002/jia2.25744>

Glidden, D. V., Mulligan, K., McMahan, V., Anderson, P. L., Guanira, J., Chariyalertsak, S., Buchbinder, S. P., Bekker, L.-G., Schechter, M., Grinsztejn, B., & Grant, R. M. (2018). Metabolic effects of preexposure prophylaxis with coformulated tenofovir disoproxil fumarate and emtricitabine. *Clinical Infectious Diseases*, 67(3), 411–419. <https://doi.org/10.1093/cid/ciy083>

Glidden, D. V., Stirrup, O. T., & Dunn, D. T. (2020). A Bayesian averted infection framework for PrEP trials with low numbers of HIV infections: Application to the results of the DISCOVER trial. *The Lancet. HIV*, 7(11), e791–e796. [https://doi.org/10.1016/S2352-3018\(20\)30192-2](https://doi.org/10.1016/S2352-3018(20)30192-2)

Gould Chadwick, E., & Edozie Ezeanolue, E. (2020). Evaluation and management of the infant exposed to HIV in the United States. *Pediatrics*, 146(5), 1–14. <https://doi.org/10.1542/peds.2020-029058>

Drugs.com. (n.d.). *GSI 225*. Retrieved March 11, 2022 from <https://www.drugs.com/imprints/gsi-225-24350.html>.

Harrison, S. E., Paton, M., Muessig, K. E., Vecchio, A. C., Hanson, L. A., & Hightow-Weidman, L. B. (2022). “Do I want PrEP or do I want a roof?”: Social determinants of health and HIV prevention in the southern United States. *AIDS Care*, 1–8. <https://doi.org/10.1080/09540121.2022.2029816>

Hartman, S., & Smith, K. E. (2020). HIV: What family medicine clinicians need to know. *Family Doctor*, 8(3), 34–37.

Hojilla, J. C., Hurley, L. B., Marcus, J. L., Silverberg, M. J., Skarbinski, J., Satre, D. D., & Volk, J. E. (2021). Characterization of HIV preexposure prophylaxis use behaviors and HIV incidence among US adults in an integrated health care system. *JAMA Network Open*, 4(8), e2122692. <https://doi.org/10.1001/jamanetworkopen.2021.22692>

Hong, C., Horvath, K. J., Stephenson, R., Nelson, K. M., Petroll, A. E., Walsh, J. L., & John, S. A. (2022). PrEP use and persistence among young sexual minority men 17–24 years old during the COVID-19 pandemic. *AIDS & Behavior*, 26(3), 631–638. <https://doi.org/10.1007/s10461-021-03423-5>



- Hoover, K. W., Huang, Y.-L. A., Tanner, M. L., Weiming Zhu, Gathua, N. W., Pitasi, M. A., DiNunno, E. A., Nair, S., & Delaney, K. P. (2020). HIV testing trends at visits to physician offices, community health centers, and emergency departments - United States, 2009-2017. *MMWR: Morbidity & Mortality Weekly Report*, 69(25), 776–780. <https://doi.org/10.15585/mmwr.mm6925a2>
- Hsu, K. K., & Rakhmanina, N. Y. (2022). Adolescents and young adults: The pediatrician's role in HIV testing and pre- and postexposure HIV prophylaxis. *Pediatrics*, 149(1), 1–18. <https://doi.org/10.1542/peds.2021-055207>
- Huang, Y., Seaton, K. E., Casapia, M., Polakowski, L., De Rosa, S. C., Cohen, K., Yu, C., Elizaga, M., Paez, C., Miner, M. D., Kelley, C. F., Maenza, J., Keefer, M., Lama, J. R., Sobieszczyk, M., Buchbinder, S., Baden, L. R., Lee, C., Gulati, V., & Sinangil, F. (2021). AIDS-VAX protein boost improves breadth and magnitude of vaccine-induced HIV-1 envelope-specific responses after a 7-year rest period. *Vaccine*, 39(33), 4641–4650. <https://doi.org/10.1016/j.vaccine.2021.06.066>
- Huang, Y.-L. A., Tao, G., Smith, D. K., & Hoover, K. W. (2021). Persistence with human immunodeficiency virus pre-exposure prophylaxis in the United States, 2012–2017. *Clinical Infectious Diseases*, 72(3), 379–385. <https://doi.org/10.1093/cid/ciaa037>
- Kanj, A., Samhoury, B., Abdallah, N., Chehab, O., & Baqir, M. (2021). Host factors and outcomes in hospitalizations for pneumocystis jirovecii pneumonia in the United States. *Mayo Clinic Proceedings*, 96(2), 400–407. <https://doi.org/10.1016/j.mayocp.2020.07.029>
- Kanny, D., Jeffries IV, W. L., Chapin-Bardales, J., Denning, P., Cha, S., Finlayson, T., Wejnert, C., & Jeffries, W. L. 4th. (2019). Racial/ethnic disparities in HIV preexposure prophylaxis among men who have sex with men - 23 urban areas, 2017. *MMWR: Morbidity & Mortality Weekly Report*, 68(37), 801–806. <https://doi.org/10.15585/mmwr.mm6837a2>
- Klein, A., & Golub, S. A. (2020). Enhancing gender-affirming provider communication to increase health care access and utilization among transgender men and trans-masculine non-binary individuals. *LGBT Health*, 7(6), 292–304. <https://doi.org/10.1089/lgbt.2019.0294>
- Klein, P. W., Geiger, T., Chavis, N. S., Cohen, S. M., Ofori, A. B., Umali, K. T., & Hauck, H. (2020). The Health Resources and Services Administration's Ryan White HIV/AIDS program in rural areas of the United States: Geographic distribution, provider characteristics, and clinical outcomes. *PloS One*, 15(3), e0230121. <https://doi.org/10.1371/journal.pone.0230121>
- Labuda, S. M., Huo, Y., Kacanek, D., Patel, K., Huybrechts, K., Jao, J., Smith, C., Hernandez-Diaz, S., Scott, G., Burchett, S., Kakkar, F., Chadwick, E. G., Dyke, R. B. V., & Study, P. H. C. (2020). Rates of hospitalization and infection-related hospitalization among human immunodeficiency virus (HIV)-exposed uninfected children compared to HIV-unexposed uninfected children in the United States, 2007–2016. *Clinical Infectious Diseases*, 71(2), 332–339. <https://doi.org/10.1093/cid/ciz820>
- Leddy, A. M., Sheira, L. A., Tamraz, B., Sykes, C., Kashuba, A. D. M., Wilson, T. E., Adedimeji, A., Merenstein, D., Cohen, M. H., Wentz, E. L., Adimora, A. A., Ofotokun, I., Metsch, L. R., Turan, J. M., Bacchetti, P., & Weiser, S. D. (2020). Food insecurity is associated with lower levels of antiretroviral drug concentrations in hair among a cohort of women living with human immunodeficiency virus in the United States. *Clinical Infectious Diseases*, 71(6), 1517–1523. <https://doi.org/10.1093/cid/ciz1007>



- Markowitz, M., Grossman, H., Anderson, P. L., Grant, R., Gandhi, M., Horng, H., & Mohri, H. (2017). Newly acquired infection with multidrug-resistant HIV-1 in a patient adherent to preexposure prophylaxis. *Journal of Acquired Immune Deficiency Syndromes (1999)*, 76(4), e104–e106. <https://doi.org/10.1097/QAI.0000000000001534>
- Mauck, D. E., Fennie, K. P., Ibañez, G. E., Fenkl, E. A., Sheehan, D. M., Maddox, L. M., Spencer, E. C., & Trepka, M. J. (2020). Estimating the size of HIV-negative MSM population that would benefit from pre-exposure prophylaxis in Florida. *Annals of Epidemiology*, 44, 52–56. <https://doi.org/10.1016/j.annepidem.2020.02.003>
- Mayer, K. H., Molina, J.-M., Thompson, M. A., Anderson, P. L., Mounzer, K. C., De Wet, J. J., DeJesus, E., Jessen, H., Grant, R. M., Ruane, P. J., Wong, P., Ebrahimi, R., Zhong, L., Mathias, A., Callebaut, C., Collins, S. E., Das, M., McCallister, S., Brainard, D. M., & Brinson, C. (2020). Emtricitabine and tenofovir alafenamide vs emtricitabine and tenofovir disoproxil fumarate for HIV pre-exposure prophylaxis (DISCOVER): Primary results from a randomised, double-blind, multicentre, active-controlled, phase 3, non-inferiority trial. *Lancet*, 396(10246), 239–254. [https://doi.org/10.1016/S0140-6736\(20\)31065-5](https://doi.org/10.1016/S0140-6736(20)31065-5)
- McComsey, G. A., Lingohr-Smith, M., Rogers, R., Lin, J., & Donga, P. (2021). Real-world adherence to antiretroviral therapy among HIV-1 patients across the United States. *Advances in Therapy*, 38(9), 4961–4974. <https://doi.org/10.1007/s12325-021-01883-8>
- McCormack, S., Dunn, D. T., Desai, M., Dolling, D. I., Gafos, M., Gilson, R., Sullivan, A. K., Clarke, A., Reeves, I., Schembri, G., Mackie, N., Bowman, C., Lacey, C. J., Apea, V., Brady, M., Fox, J., Taylor, S., Antonucci, S., Khoo, S. H., & Rooney, J. (2016). Pre-exposure prophylaxis to prevent the acquisition of HIV-1 infection (PROUD): Effectiveness results from the pilot phase of a pragmatic open-label randomised trial. *Lancet*, 387(10013), 53–60. [https://doi.org/10.1016/S0140-6736\(15\)00056-2](https://doi.org/10.1016/S0140-6736(15)00056-2)
- Mulligan, K., Glidden, D. V., Anderson, P. L., Liu, A., McMahan, V., Gonzales, P., Ramirez-Cardich, M. E., Namwongprom, S., Chodacki, P., de Mendonca, L. M. C., Wang, F., Lama, J. R., Charialertsak, S., Guanira, J. V., Buchbinder, S., Bekker, L.-G., Schechter, M., Veloso, V. G., & Grant, R. M. (2015). Effects of emtricitabine/tenofovir on bone mineral density in hiv-negative persons in a randomized, double-blind, placebo-controlled trial. *Clinical Infectious Diseases : An Official Publication of the Infectious Diseases Society of America*, 61(4), 572–580. <https://doi.org/10.1093/cid/civ324>
- Nesheim, S. R., Balaji, A., Hu, X., Lampe, M., & Dominguez, K. L. (2021). Opportunistic illnesses in children with HIV infection in the United States, 1997-2016. *The Pediatric Infectious Disease Journal*, 40(7), 645–648. <https://doi.org/10.1097/INF.0000000000003154>
- Olakunde, B. O., Pharr, J. R., & Adeyinka, D. A. (2020). HIV testing among pregnant women with prenatal care in the United States: An analysis of the 2011-2017 national survey of family growth. *International Journal of STD & AIDS*, 31(7), 680–688. <https://doi.org/10.1177/0956462420921715>
- Pampati, S., Emrick, K., Siegler, A. J., & Jones, J. (2021). Changes in sexual behavior, PrEP adherence, and access to sexual health services because of the COVID-19 pandemic among a cohort of PrEP-using MSM in the South. *Journal of Acquired Immune Deficiency Syndromes (1999)*, 87(1), 639–643. <https://doi.org/10.1097/QAI.0000000000002640>



- Peruski, A. H., Wesolowski, L. G., Delaney, K. P., Chavez, P. R., Owen, S. M., Granade, T. C., Sullivan, V., Switzer, W. M., Xueyuan Dong, Brooks, J. T., Joyce, M. P., & Dong, X. (2020). Trends in HIV-2 diagnoses and use of the HIV-1/HIV-2 differentiation test - United States, 2010-2017. *MMWR: Morbidity & Mortality Weekly Report*, 69(3), 63–66. <https://doi.org/10.15585/mmwr.mm6903a2>
- Saá, P., Townsend, R. L., Wells, P., Janzen, M. A., Brodsky, J. P., & Stramer, S. L. (2020). Qualification of the Geenius HIV 1/2 supplemental assay for use in the HIV blood donation screening algorithm. *Transfusion*, 60(8), 1804–1810. <https://doi.org/10.1111/trf.15819>
- San Francisco AIDS Foundation. (2019, November). *Side-by-side comparison: Truvada and Descovy for PrEP*. <https://www.sfaf.org/resource-library/side-by-side-comparison-truvada-and-descovy-for-prep/>
- Seelman, K. L., Kattari, S. K., Harvey, P., & Bakko, M. (2020). Trans men’s access to knowledgeable providers and their experiences in health care settings: Differences by demographics, mental health, and degree of being “out” to providers. *Health & Social Work*, 45(4), 229–239. <https://doi.org/10.1093/hsw/hlaa030>
- Sharma, A., Paredes-Vincent, A., & Kahle, E. M. (2021). Awareness, Utilization, and Preferences for Traditional and contemporary HIV prevention strategies among Facebook and Instagram-using MSM in the United States. *Journal of the International Association of Providers of AIDS Care*, 1–17. <https://doi.org/10.1177/23259582211024770>
- Solomon, M. M., Lama, J. R., Glidden, D. V., Mulligan, K., McMahan, V., Liu, A. Y., Guanira, J. V., Veloso, V. G., Mayer, K. H., Chariyalertsak, S., Schechter, M., Bekker, L.-G., Kallás, E. G., Burns, D. N., & Grant, R. M. (2014). Changes in renal function associated with oral emtricitabine/tenofovir disoproxil fumarate use for HIV pre-exposure prophylaxis. *AIDS*, 28(6), 851–859. <https://doi.org/10.1097/QAD.000000000000156>
- Struble, C. A., Bauer, S. J., Lundahl, L. H., Ghosh, S., & Ledgerwood, D. M. (2022). Electronic cigarette use among sexual minority and heterosexual young adults in a U.S. national sample: Exploring the modifying effects of advertisement exposure. *Preventive Medicine*, 155, 106926. <https://doi.org/10.1016/j.ypmed.2021.106926>
- Stults, C. B., Grov, C., Anastos, K., Kelvin, E. A., & Patel, V. V. (2020). Characteristics associated with trust in and disclosure of sexual behavior to primary care providers among gay, bisexual, and other men who have sex with men in the United States. *LGBT Health*, 7(4), 208–213. <https://doi.org/10.1089/lgbt.2019.0214>
- Sullivan, P. S., Sanchez, T. H., Zlotorzynska, M., Chandler, C. J., Sineath, R. C., Kahle, E., & Tregear, S. (2020). National trends in HIV pre-exposure prophylaxis awareness, willingness and use among United States men who have sex with men recruited online, 2013 through 2017. *Journal of the International AIDS Society*, 23(3), e25461. <https://doi.org/10.1002/jia2.25461>
- Tang, E. C., Vittinghoff, E., Philip, S. S., Doblecki-Lewis, S., Bacon, O., Chege, W., Coleman, M. E., Elion, R., Buchbinder, S., Kolber, M. A., Liu, A. Y., & Cohen, S. E. (2020). Quarterly screening optimizes detection of sexually transmitted infections when prescribing HIV preexposure prophylaxis. *AIDS (London, England)*, 34(8), 1181–1186. <https://doi.org/10.1097/QAD.0000000000002522>



- Tassiopoulos, K., Huo, Y., Patel, K., Kacanek, D., Allison, S., Siminski, S., Nichols, S. L., Mellins, C. A., & PHACS), Pediatric HIV/AIDS Cohort Study. (2020). Healthcare transition outcomes among young adults with perinatally acquired human immunodeficiency virus infection in the United States. *Clinical Infectious Diseases*, 71(1), 133–141. <https://doi.org/10.1093/cid/ciz747>
- van der Laan, L. E., Garcia-Prats, A. J., Schaaf, H. S., Winckler, J. L., Draper, H., Norman, J., Wiesner, L., McIlleron, H., Denti, P., & Hesselning, A. C. (2021). Pharmacokinetics and drug-drug interactions of abacavir and lamuvudine co-administered with antituberculosis drugs in HIV-positive children treated for multidrug-resistant tuberculosis. *Frontiers in Pharmacology*, 12, 722204. <https://doi.org/10.3389/fphar.2021.722204>
- Vézina, D., Shang Yu Gong, Tolbert, W. D., Shilei Ding, Dung Nguyen, Richard, J., Gendron-Lepage, G., Melillo, B., Smith III, A. B., Pazgier, M., & Finzi, A. (2021). Stabilizing the HIV-1 Envelope Glycoprotein State 2A Conformation. *Journal of Virology*, 95(5), 1–11. <https://doi.org/10.1128/JVI.01620-20>
- Vijayan, V., Naeem, F., & Veesenmeyer, A. F. (2021). Management of infants born to mothers with HIV infection. *American Family Physician*, 104(1), 58–62.
- Wahnich, A., Gandhi, A. D., Cleghorn, E., Estacio, K., Blackstock, O. J., Myers, J. E., Abraham, B., & Edelstein, Z. R. (2021). Public health detailing to promote HIV pre- and postexposure prophylaxis among women’s healthcare providers in New York City. *American Journal of Preventive Medicine*, 61, S98–S107. <https://doi.org/10.1016/j.amepre.2021.05.032>
- Walters, S. M., Coston, B., Neaigus, A., Rivera, A. V., Starbuck, L., Ramirez, V., Reilly, K. H., & Braunstein, S. L. (2020). The role of syringe exchange programs and sexual identity in awareness of pre-exposure prophylaxis (PrEP) for male persons who inject drugs. *International Journal of Drug Policy*, 77, Article 102671. <https://doi.org/10.1016/j.drugpo.2020.102671>
- Watson, R. J., Eaton, L. A., Maksut, J. L., Rucinski, K. B., & Earnshaw, V. A. (2020). Links between sexual orientation and disclosure among black MSM: Sexual orientation and disclosure matter for prep awareness. *AIDS and Behavior*, 24(1), 39–44. <https://doi.org/10.1007/s10461-019-02696-1>
- Weiss, G., Smith, D. K., Newman, S., Wiener, J., Kitlas, A., & Hoover, K. W. (2018). PrEP implementation by local health departments in US cities and counties: Findings from a 2015 assessment of local health departments. *PLoS One*, 13(7), e0200338. <https://doi.org/10.1371/journal.pone.0200338>
- Wohl, D., Ruane, P., Hosek, S., Creticos, C., Morris, S., Phoenix, J., Ramgopal, M., Brinson, C., Tremblay, C., Carter, C. C., Wong, P., Brainard, D. M., McCallister, S., Das, M., & Thompson, M. A. (2019). 1288. Bone safety outcomes with F/TAF vs. F/TDF for PrEP in the DISCOVER Trial. *Open Forum Infectious Diseases*, 6(Suppl 2), S464. <https://doi.org/10.1093/ofid/ofz360.1151>
- Yusuf, H., Fields, E., Arrington-Sanders, R., Griffith, D., & Agwu, A. L. (2020). HIV preexposure prophylaxis among adolescents in the US: A Review. *JAMA Pediatrics*, 174(11), 1102–1108. <https://doi.org/10.1001/jamapediatrics.2020.0824>



Image References

Slide(s)	Image designation/description/position	Attribution
2	Disclosures	Image licensed through Creative Commons: https://www.picpedia.org/real-estate/d/disclosures.html
4-11	CDC Report Figure A Figure 1 Figure 2 Figure 3 Figure 5 Figure 6 Figure 13 Figure 22	Source: https://www.cdc.gov/hiv/pdf/library/reports/surveillance/cdc-hiv-surveillance-report-2020-updated-vol-33.pdf
12	HIV Replication Cycle	Image licensed through Creative Commons by NIH NIAID: https://www.niaid.nih.gov/diseases-conditions/hiv-replication-cycle
19	IV Drug Injection Image	Image licensed through Creative Commons by PXhere: https://pxhere.com/en/photo/774066
21	Pregnancy Image	Image licensed through Creative Commons by StockSnap: https://stocksnap.io/photo/pregnant-woman-40B226DC63
24	Child Hand Halt	Image licensed through Creative Commons by Wallpaper Flare: https://www.wallpaperflare.com/the-hand-child-cover-up-do-not-do-not-shoot-the-prohibition-of-wallpaper-atdyk
25	OraQuick Testing	Image licensed through Creative Commons by Wikimedia Commons: https://commons.wikimedia.org/wiki/File:Oraquick.jpg
33	Post-Exposure Risk	Source: https://stacks.cdc.gov/view/cdc/38856
34	Post-Exposure Algorithm	Source: https://stacks.cdc.gov/view/cdc/38856
35	Post-Exposure Rx	Source: https://stacks.cdc.gov/view/cdc/38856
36	Post-Exposure Labs	Source: https://stacks.cdc.gov/view/cdc/38856
37	Risk Gauge	Image licensed through Creative Commons by PXhere: https://pxhere.com/en/photo/1444649
38	Pre-Exposure Prophylaxis	Source: https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2021.pdf
39	Condoms	Image licensed through Creative Commons by Pixabay https://pixabay.com/photos/condoms-condom-rubber-prevention-3112008/
40	Oral PrEP Labs	Source: https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2021.pdf
41	"2-1-1" PrEP Regimen	Source: https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2021.pdf
42	Money	Image licensed through Creative Commons by flickr: https://www.flickr.com/photos/tracy_olson/61056391
46	AANP	Image licensed through Creative Commons by Wikimedia Commons: https://commons.wikimedia.org/wiki/File:American_Association_of_Nurse_Practitioners_logo_031918.png
47	Moving Forward	Image licensed through Creative Commons by Pixy: https://pixy.org/1601486/
48	Thank You	Image licensed through Creative Commons: https://publicdomainvectors.org/en/free-clipart/Thank-You-note-vector-drawing/25496.html



Pharmacologic and Non-Pharmacologic Methods in the Prevention of HIV: 2023

Christopher W. Blackwell, Ph.D., APRN,
ANP-BC, AGACNP-BC, CNE, FAANP,
FAAN

*Associate Professor & Program Director
AGACNP Programs, College of Nursing
Academic Health Sciences Center
University of Central Florida
Orlando, Florida*

